(REV 10-94)	PARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE					
TRANSMITTAL LETTER		13390.2USWO				
DESIGNATED/ELECTE						
CONCERNING A FILIN	G UNDER 35 U.S.C. 371					
		UNKNOWN 0 / 018719				
	INTERNATIONAL FILING DATE	PRIORITY DATE CLAIMED				
PCT/EP00/05517	JUNE 15, 2000	JUNE 17, 1999				
TITLE OF INVENTION						
USE OF GROWTH HORMONE (HGH) FO	OR THE TREATMENT OF SEXUAL FUNC	CTIONAL DISTURBANCES				
APPLICANT(S) FOR DO/EO/US						
BECKER, Armin Johannes; STIEF, Christia	ian George; UCKERT, Stefan; JONAS, Udo					
Applicant herewith submits to the United States De	signated/Elected Office (DO/EO/US) the following	g items and other information:				
[X] This is a <b>FIRST</b> submission of items cond	corning a filma under 35 U.S.C. 371					
2. [ ] This is a <b>SECOND</b> or <b>SUBSEQUEN</b>	NT submission of items concerning a filing under 3	35 U.S.C. 371.				
<ol><li>[X] This express request to begin national exa</li></ol>	amination procedures (35 U.S.C. 371(f)) at any time	e rather than delay				
examination until the expiration of the appear [X] A proper Demand for International Prelim	plicable time limit set in 35 U.S.C. 371(b) and PCT ninary Examination was made by the 19th month fr	l' Articles 22 and 39(I).  rom the earliest claimed priority date.				
		on the carries carries proving access				
[X] A copy of the International Application as a. [X] is transmitted herewith (required	s filed (35 U.S.C. 371(c)(2)) I only if not transmitted by the International Bureau					
b. [X] has been transmitted by the Inter	rnational Bureau.					
c. [ ] is not required, as the application was filed in the United States Receiving Office (RO/US) [ ] A translation of the International Application into English (35 U.S.C. 371(c)(2)).						
nana nana nana nana nana nana nana nan						
[X] Amendments to the claims of the Internati	ional Application under PCT Article 19 (35 U.S.C.	. 371(c)(3))				
c. [ ] have not been made; however	er, the time limit for making such amendments has	s NOT expired.				
d. [X] have not been made and will not	be made.					
N I I I I I I I I I I I I I I I I I I I	ne claims under PCT Article 19 (35 U.S.C. 371(c)(3	3)).				
[X] An unsigned oath or declaration of the inv	entor(s) (35 U.S.C. 371 (c)(4)).					
0. [ ] A translation of the annexes to the Int (35 U.S.C. 371(c)(5)).	ternational Preliminary Examination Report under I	PCT Article 36				
tems 11. to 16. below concern document(s) or int						
1. [ ] An Information Disclosure Statement	under 37 CFR 1.97 and 1.98.	İ				
2. [ ] An assignment document for recording	An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.					
[X] A FIRST preliminary amendment.     [ ] A SECOND of SUBSEQUENT preliminary amendment.	minary amendment.					
4. [ ] A substitute specification.						
. [ ] A change of power of attorney and/or address letter.						
[X] Other items or information: PCT/EP00/05517, PCT/ISA/210, PCT/IB/308; PCT/IPEA/409						

U.S. APPLICATION NO (If know		INTERNATIONAL APPLICATION N	NO	ATTORNEY'S DOCKET NUMBER		
UNKNOWN 10/01871 9CT/EP00/05517 13390.2USV						
17. [X] The following fees are submitted:			CALCULATIONS P	PTO USE ONLY		
BASIC NATIONAL FEE (37 CFR 1.492(a) (1)-(5)):						
		or JPO	\$890.00			
	ninary examination fee paid	to USPTO	\$710.00			
		oaid to USPTO (37 CFR 1.48 7 CFR 1.445(a)(2))				
	al preliminary examination f h fee (37 CFR 1.445(a)(3)) p	fee (37 CFR 1.482) nor paid to USPTO	\$1040.00			
	ninary examination fee paid sefied provisions of PCT Artic	to USPTO (37 CFR 1.482)	\$100.00			
	ENTER APPROF	PRIATE BASIC FEE	AMOUNT =	\$890.00		
	or furnishing the oath or decl t claimed priority date (37 C	claration later than [] 20 [] 3 CFR 1.492(e)).	50	\$0		
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE			
Total claims	11 -20 =	0	X \$18.00	\$0		
findependent claims	2 -3 =	0	X \$84.00	\$0		
MULTIPLE DEPENDE	ENT CLAIM(S) (if applicabl	le)	+ \$260.00	\$0		
	TOTAL	OF ABOVE CALCU	LATIONS =	\$890.00		
	ing by small entity, if applica	cable. Small entity status is o				
pursuant to 37 CFR 1.27	7			\$0		
SECOND SE			UBTOTAL =	\$890.00		
Processing fee of \$130.0 months from the earliest	00 for furnishing the English t claimed priority date (37 Cl	n translation later than [] 20 FR 1.492(f).	[]30	\$0		
agency.		TOTAL NATIO	NAL FEE =	\$890.00		
Eee for recording the enc	closed assignment (37 CFR copriate cover sheet (37 CFR	1.21(h)). The assignment m R 3.28, 3.31). \$40.00 per pro	nust be	\$0		
11	Spring over and a	TOTAL FEES EN		\$890.00		
		TOTAL PEEG EA	(CLUSED –			
				Amount to be: refunded	\$0	
77 Cl 1() in the	22222.00			charged	\$0	
a. [X] Check(s) in the	e amount of \$890.00 to cover	r the above fees is enclosed.			_	
b. [] Please charge my Deposit Account No in the amount of \$ to cover the above fees.  A duplicate copy of this sheet is enclosed.						
<ul> <li>[X] The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 13-2725.</li> </ul>						
NOTE: Where an appr 1.137(a) or (b)) must be	ropriate time limit under 3 e filed and granted to resto	37 CFR 1.494 or 1.495 has a preting the application to pendi	not been met, a ping status.	petition to revive (37 CFR	t	
SEND ALL CORRESPONDENCE	то			2 /		
Ronald A. Daignault  MERCHANT & GOULD  P.O. Box 2903  SIGNATURE: Acade A						
P.O. Box 2903						
Minneapolis, MN 554	102-0903		NAN	ME: Ronald A. Daignault		
		GISTRATION NUMBER	25,968			

Applicant:

BECKER, et al.

Docket:

13390.2USWO

Title:

USE OF GROWTH HORMONE (HGH) FOR THE TREATMENT OF SEXUAL

FUNCTIONAL DISTURBANCES

#### CERTIFICATE UNDER 37 CFR 1 10

'Express Mail' mailing label number EL669945412US

Date of Deposit December 14, 2001

I hereby certify that this paper or fee is being deposited with the United States Postal Service 'Express Mail Post Office To Addressee' service under 37 CFR 1 10 and is addressed to the Commissioner for Patents, Washington, D.C. 20231

**BOX PATENT APPLICATION** 

Commissioner for Patents Washington, D.C. 20231

Sir:

We are transmitting herewith the attached:

Transmittal sheet, in duplicate, containing Certificate under 37 CFR 1.10.

National Stage PCT Patent Application: Spec. 8 pgs; 11 claims; Abstract 1 pgs.

The fee has been calculated as shown below in the 'Claims as Filed' table.

2 sheets of formal drawings

An unsigned Combined Declaration and Power of Attorney

A check in the amount of \$890.00 to cover the Filing Fee

Other: PCT/EP00/05517, Preliminary Amendment, PCT/ISA/210; PCT/IB/308; PCT/IPEA/409

Return postcard

#### CLAIMS AS FILED

		CLAIMS F	70 L IT	ED				
Number of Claims Filed		In Excess of:		Number Extra		Rate		Fee
Basic Filing Fee								\$890.00
Total Claims								
11	-	20	=	0	X	18.00	=	\$0.00
Independent Claims								
2	-	3	=	0	X	84.00	=	\$0.00
MULTIPLE DEPENDENT C	CLAIM I	FEE						\$0.00
TOTAL FILING FEE								\$890.00

Please charge any additional fees or credit overpayment to Deposit Account No. 13-2725. A duplicate of this sheet is enclosed.

MERCHANT & GOULD P.C.

P.O. Box 2903, Minneapolis, MN 55402-0903

(612) 332-5300

Name: Ronald A. Daignault

Reg. No. 25,968

Initials: RADaignault/rw

531 7000 70000

4DEC 2001

# IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant:

Becker, et al.

Docket No.:

13390.2USWO

Serial No.:

unknown

Filed:

concurrent herewith

Int'l Appln No.:

PCTEP00005517

Int'l Filing Date:

June 15, 2000

Title:

USE OF GROWTH HORMONE (HGH) FOR THE TREATMENT OF

SEXUAL FUNCTIONAL DISTURBANCES

CERTIFICATE UNDER 37 CFR 1.10

'Express Mail' mailing label number: EL669945412US

Date of Deposit: December 14, 2001

I hereby certify that this paper or fee is being deposited with the United States Postal Service 'Express Mail Post Office To Addressee' service under 37 CFR 1.10 and is addressed to the Commissioner for Patents, Washington, D.C. 20231.

By: Name: Chris Stordahl

## PRELIMINARY AMENDMENT

**Box PCT** 

**Assistant Commissioner for Patents** 

Washington, D. C. 20231

Dear Sir:

In connection with the above-identified application filed herewith, please enter the following preliminary amendment:

## IN THE ABSTRACT

Insert the attached Abstract page into the application as the last page thereof.

A courtesy copy of the present specification is enclosed herewith. However, the World Intellectual Property Office (WIPO) copy should be relied upon if it is already in the U.S. Patent Office.

#### REMARKS

A new abstract page is supplied to conform to that appearing on the publication page of the WIPO application, but the new Abstract is typed on a separate page as required by U.S. practice.

Applicants respectfully request that the preliminary amendment described herein be entered into the record prior to calculation of the filing fee and prior to examination and consideration of the above-identified application.

If a telephone conference would be helpful in resolving any issues concerning this communication, please contact Applicants' primary attorney-of record, Ronald A. Daignault (Reg. No. 25,968), at (612) 371.5381.

Respectfully submitted,

MERCHANT & GOULD P.C.

P.O. Box 2903

Minneapolis, Minnesota 55402-0903

(612) 332-5300

Dated: December 14, 2001

RAD/rw

Ronald A. Dalgnault Reg. No. 25,968

### **ABSTRACT** PCT/EP00/05517

# USE OF GROWTH HORMONE (HGH) FOR THE TREATMENT OF SEXUAL **FUNCTIONAL DISTURBANCES**

The invention relates to the use of the human growth hormone (hGH, GH) either on its own or in combination with active substances which result in GH stimulation, have an effect which is analogous to GH or which promote the release of IGF-I in the production of medicaments which are used to treat sexual functional disturbances in both sexes. The invention also relates to a method for treating the above-mentioned dysfunctions.

CERTIFICATE UNDER 37 CFR 1.10

'Express Mail' mailing label number: EL669945412US

Date of Deposit: December 14, 2001

I hereby certify that this paper or fee is being deposited with the United States Postal Service 'Express Mail Post Office To Addressee' service under 37 CFR 1.10 and is addressed to the Commissioner for Patents, Washington, D.C. 20231.

Name: Chris Stordahl

Rec'd PCT/FTO 21 MAR 2002

10/018719 ENT

S/N 10/018719

# IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant:

Becker, et al.

Docket No.:

13390.2USWO

Serial No.:

10/018719

Filed:

December 14, 2001

Int'l Appln No.:

PCTEP00005517

Int'l Filing Date:

June 15, 2000

Title:

USE OF GROWTH HORMONE (HGH) FOR THE TREATMENT OF

SEXUAL FUNCTIONAL DISTURBANCES

CERTIFICATE UNDER 37 CFR 1.10

'Express Mail' mailing label number: EV072823491

Date of Deposit: March 21, 2002

I hereby certify that this paper or fee is being deposited with the United States Postal Service 'Express Mail Post Office To Addressee' service under 37 CFR 1.10 on the date indicated above and is addressed to the Commissioner of Patents and Trademarks, Washington, D.C. 20231.

Chris Stordahl

# PRELIMINARY AMENDMENT

Box PCT Assistant Commissioner for Patents Washington, D. C. 20231

Dear Sir:

In connection with the above-identified application filed herewith, please enter the following preliminary amendment.

The present application was filed in the GERMAN language. Applicant now supplies the English translation of the application and the verified English translation of the specification are supplied herewith.

## IN THE SPECIFICATION

Please amend the specification as follows:

# Administration of Growth Hormone (hGH) for Therapy of Sexual Functional Disorders

#### **Technical Field**

[001] This invention concerns the introduction of human growth hormone (hGH, GH) for the manufacture of medicaments for the treatment of sexual functional disorders in both male and female patients as well as the specific methods of treatment undertaken.

# **Background of the Invention**

[002] Symptoms indicating sexual functional disorders are, for example, lack or loss of libido, problems relating to orgasms, insufficient lubrication and erectile dysfunction (ED). It could be deduced from certain cases that the basic aetiologies leading to these sexual functional disorders were due to a number of different reasons. Apart from the cases of mixed aetiologies, vascular (arterial, venous), psychogenic, neurogenic, medicamentous-induced and cavernous sexual functional disorders are also differentiated.

[003] Taking into consideration the nature of the underlying aetiologies, causal therapy of the sexual functional disorders is undertaken whenever possible. Up till now this method of therapy has only proved successful in the rare cases (e.g. by psychotherapy, hormone treatment, a change-over of medication) so that the main methods of therapy still remain unspecific.

[004] There are many different methods of therapy available for males with ED compared to those available for females with sexual functional disorders. These include oral, topical, intracavernous, intraurethral and also a combination of drugs. These methods do not constitute a causal therapy, rather the aim is to achieve a direct or indirect relaxation (flaccidity) of the corpus cavernosum smooth musculature and the penile arteries. Together with an increase of blood circulation, penile erection is achieved. Furthermore, the vacuum pump, arterial shunt procedure, venous closure operations and penile prosthesis implantations are also methods used for therapy. Until the introduction of sildenafil (VIAGRA®), the most widely used form of therapy involved the administration of intracavernous vasoactive substances. At the present time, sildenafil is used

as the so-called "first line therapy" providing there are no known contraindications. The oral phosphodiesterase type 5 inhibitor (PDE5) does not provide a basis for causal therapy. With the inhibition of PDE5, hydrolysis of cyclic guanosinmonophosphate (cGMP) of an intracellular second messenger is prohibited, resulting in relaxation of the corpus cavernosum smooth musculature. This effective mechanism is assumed beneficial for the increase of lubrication in women, however, recent studies have yet to prove its effectivity.

## **Summary of the Invention**

[005] The aim of this innovative breakthrough was to present a new therapy for both males and females suffering from sexual functional disorders.

[006] Surprisingly, it has been shown that the growth hormone (hGH) plays an essential role in sexual stimulation, as an enormous unexpected increase of this hormone was seen to be present at the onset of sexual stimulation.

[007] The focal point of this breakthrough is therefore the use of hGH for the manufacture of medicaments for the treatment of sexual functional disorders in both males and females with e.g. lack or loss of libido, problems relating to orgasms, insufficient lubrication and erectile dysfunction and for the therapy of the aforesaid functional disorders.

[008] Accordingly, the present invention is a method of treating female and male sexual functional disorders that includes administering an effective amount of growth hormone. The sexual functional disorder is manifested by a lack or loss of libido, problems relating to orgasms, insufficient lubrication, or erectile dysfunction; there is an insufficient increase in hGH concentration occurs during sexual stimulation or a hGH deficiency exists. A further aspect is the use of hGH for the therapy of functional disorders in synergic combination with effective substances which result in GH stimulation, induce a GH analogous effect or promote IGF-I release.

# **Brief Description of the Drawings**

[010] **Figure 1** shows the average values and standard deviations of the growth hormone (hGH) concentrations (ng/ml) in cavernous and peripheral blood samples taken from 35 healthy probands during the four different phases of the penile erectile tissue (flaccidity, tumescence, rigidity and detumescence).

[011] **Figure 2** shows the average values and standard deviations of the growth hormone (hGH) concentrations (ng/ml) in the cavernous and peripheral blood samples during the three different penile phases (flaccidity, tumescence and detumescence) in 36 patients with erectile dysfunction. Rigidity was not achieved due to disorder of patient. The axis scale was selected as in Figure 1 in order to demonstrate clearly the difference (P<0.05).

[012] **Figure 3** shows the average values and standard deviations of the dose-dependent decrease in relaxation of 12 human corpus cavernosum strips after application of recombinant hGH.

[013] **Figure 4** shows the average values and standard deviations of dose-dependent increase of cyclic guanosinmonophasphate (cGMP) of 3 human corpus cavernosum strips respectively, following incubation with recombinant hGH or sodium nitroprusside (SNP). Incubation with SNP was carried out using a concentration of 0.01 and 1 µMol. For this reason, no value for SNP was achieved with 0.0001 µMol.

## **Detailed Description of the Invention**

[014] In order to gain a better understanding of the physiology of a naturally induced erection and the pathophysiology of erectile dysfunction, a new method of investigation was developed. This involved detection of endogenous human neurotransmitters, neuromodulators and hormones which might have some connection with an erection or its sexual function. These new methods of investigation aim to improve the diagnostics and therapy (application of endogenous substances and causal therapy) for those patients with sexual functional disorders.

[015] Blood was taken simultaneously from the corpus cavernosum (CC, cavernous) and the cubital vein (CV, peripheral) in 35 healthy probands during the phases of flaccidity, tumescence, rigidity and detumescence. Audiovisual and tactile means were then provided to aid sexual stimulation (Figure 1). The procedure involving 36 patients with ED was identical to that of the healthy probands with the exception of blood withdrawal during the rigidity phase, (this penile erection phase cannot be achieved in patients with ED) (Figure 2). The hGH concentrations were determined with an immunoradiometric assay (IRMA). This form of investigation resulted in several new findings.

[016] The highest increase of hGH concentration was found during tumescence, namely the point in time when sexual stimulation is at its peak. The peripheral and cavernous hGH concentrations showed no significant differences in direct comparison with all penile phases. Peripheral blood withdrawal proved sufficient. When comparing the healthy probands with the patients, there were significant differences with regards to the hGH concentrations, in particular, a significantly reduced increase of the hGH concentration during the tumescence phase.

[017] These data show for the first time the surprising causal connection of hGH formed by the hypophysis with sexual stimulation and the resulting penile erection. The reduced expression of hGH on sexual stimulation in patients is further proof for the significance of this hormone, the lack of which is connected with sexual functional disorders and erectile dysfunction in this study.

[018] By means of extensive in vitro investigations using human corpus cavernosum (CC) tissue as well as the in vivo results described, important indications could be deduced for the possible physiological connections between hGH and penile erection. Organ bath experiments (in vitro method to evaluate the relaxing properties of substances) with human CC were carried out to assess dose-dependent relaxations following application of hGH (**Figure 3**).

[019] Incubation experiments (in vitro method to evaluate the content of cyclic nucleotides in tissues in response to drug exposition, in this case cGMP, after incubation with various substances) with human CC were shown to have dose-dependent higher cGMP concentrations

after application of hGH than was the case after incubation with sodium nitroprusside (SNP), a classic NO donator (Figure 4).

[020] Based on our human findings, it can be assumed that hGH plays a decisive role in sexual function (sexual stimulation), in particular, in penile erection. Furthermore, it was shown that the peripheral reaction of hGH induced an increase of cGMP, thereby physiologically forming a link between relaxation of CC with that of the ensuing erection. Due to the anatomical similarities in the structure of the penis and the clitoris and the physiological conformities regarding sexual stimulation (e.g. congestion of the genital organs mediated by neurotransmitter on relaxation of the smooth musculature), the described reaction of hGH in males must also apply to females too, since hGH is produced by the hypophysis in both sexes, therefore the same effect must also be evident in both sexes.

[021] With reference to the effect of hGH, it is already known that hGH does not focus on any particular tissue and that the activity and metabolism (anabolic) is increased in different tissues in both men and women. The growth hormone stimulates, for example, body growth (substitution in insufficient hGH-caused hyposomia) and protein metabolism (possible indication in cachexia, severe burn injuries and also anabolic abuse). Under the influence of hGH, an insulin-like growth factor I (IGF-I) is formed mainly in the liver but also in other tissues. This polypeptide (IGF-I) plays a significant mediating role in the process induced by hGH (Merimee T.J. and Grant M.B.: Growth hormone and its disorders. In: Principles and Practice of Endocrinology and Metabolism. Edited by Becker, K.L., Philadelphia, J.B. Lippincott Company, pp. 125-134, 1990).

[022] The most recent findings in humans show that there is a systemic increase of NO (nitric oxide) and cGMP under a substitution of recombinant produced hGH (r-hGH) in patients with hGH deficiency (Böger R.H. et al.: Nitric oxide may mediate the hemodynamic effects of recombinant growth hormone in patients with acquired growth hormone deficiency. J. Clin. Invest. 98: 2706-2713, 1996). This NO-cGMP path presents a very important positive significance in achieving penile erection (Burnett A.L. et al.: Nitric oxide: a physiologic mediator or penile erection. Science 257: 905, 1993). Likewise, recent findings derived from animal experiments using rats were able to show that an increase of NOS (nitric oxide synthase)-

containing nerves (generating NO) in CC and dorsal penile nerves occurred under substitution of hGH. This took place despite initiation of neurogenic damage some weeks before (Jung G.W. et al.: Growth hormone enhances regeneration of nitric oxide synthase-containing penile nerves after cavernous neurotomy in rats. J. Urol. 160: 1899-1904, 1998). The patent WO98/42361 (Human erectile dysfunction and methods of treatment) stems from these results and describes the indication of hGH therapy for the prevention and treatment of neurogenic erectile dysfunction of different aetiologies (condition following extensive pelvic operations or pelvic trauma, diabetes, alcoholism and aging process).

- [023] Our findings on human models show for the first time a positive causal connection between sexual stimulation, increase of hGH and penile erection. The reduced (often totally lacking) increase of hGH in patients with ED emphasizes the importance of this hormone. The in vitro data conclude that the NO-cGMP path is activated by hGH, leading to relaxation of the CC, thus resulting in penile erection.
- [024] The appropriate therapy for all patients (both sexes) with sexual functional disorders includes peripheral blood sample to determine the basal hGH concentration. This form of therapy is undertaken independent of the underlying aetiology(ies). Following this, a further blood sample is taken under sexual stimulation (audiovisual, tactile) in order to detect the stimulated hGH concentration. In cases of insufficient or no reaction at all to sexual stimulation (e.g. lubrication, penile erection) and inadequate increase of hGH concentration, a continuous, strictly controlled therapy with hGH should ensue for a certain period of time (e.g. 2 6 months).
- [025] Suitable pharmaceutical preparations for therapy include solid or liquid forms of administration for oral intake, such as tablets, capsules or emulsions, parenteral forms of administration for injection or non-invasive application or transdermal topical systems, such as plasters, creams, gels, lotions or transdermal films. The administered amount for successful therapy lies between 0.01 and 500 mg per dosage unit; recommended is between 0.1 and 100 mg.

[026] Improvement of therapy outcome can be achieved by administration of a combination of medicaments containing, besides hGH, a synergic combination of substances which lead to GH stimulation, induce a GH analogous effect or promote IGF-I release.

[027] These substances do not have to be combined into one particular medication, but can be administered in separate suitable galenic preparations to be taken at the same time or taken separately according to the specific course of therapy. It is essential that the specialist instructs and informs the patient with regards to the suitable dosage or in which combination the medicaments should be taken, likewise which substance should be administered to ensure the best possible therapy outcome. Furthermore, it is permissible to combine several of the named substances to treat the individual patient accordingly.

[028] The suitable substances to be used as a combination therapy in order to achieve GH stimulation are familiar to the specialist. For example, arginine, alpha 1 and alpha 2-agonists, such as clonidine, norepinephrine or salbutamol, glucagon, pyridostigmine, galanine, GH-releasing hormone, NPY (neuropeptide Y) and dopamine agonists, such as apomorphine, quinpirole or cabergoline.

[029] Suitable substances which induce a GH analogous effect include, for instance, GHRP (growth hormone, releasing hexapeptide, hexareline), GH releasing peptide 1, 2, 6 and non-peptidergic agonists of growth hormone releasing peptide such as MK 0677, EP 51389 (2-methylalanyl-2-methyl-D-tryptophyl-2-methyl-D-tryptophanamide), L 692429 (3-amino-3-methyl-N-[(3R)-2,3,4,5-tetrahydro-2-oxo-1-[[1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-1H-1-benzazepine-3-yl]butanamide], or L 692585 (3-[[(2R)-2hydroxypropyl]amino]-3-methyl-N-[(3R)-2,3,4,5-tetrahydro-2-oxo-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-1H-1-benzazepine-3-yl]-butanamide).

[030] Suitable substances which promote IGF-I release include, for example, cannabinoide such as e.g. HU-210 (3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-6H-dibenzo[b,d]pyran-9-methanol) or serotonine receptor agonists such as e.g. 8-OH

DPAT (8-hydroxy-2-dipropylamino)tetraline), or SC 53116 (4-amino-5-chloro-N-[[(1s,7aS)-hexahydro-1H-pyrrolizine-1-yl]methyl]-2-methoxy-benzamide)

## IN THE CLAIMS

Please cancel claims 1-11 without prejudice or disclaimer. Add new claims 12-18.

- 12. (New) A method of treating femaled and male sexual functional disorders comprising administering an effective amount of a growth hormone, wherein the sexual functional disorder is manifested by lack or loss of libido, problems relating to orgasms, insufficient lubrication, or erectile dysfunction and wherein insufficient increase in hGH concentration occurs during sexual stimulation or a hGH deficiency exists.
- 13. (New) The method of claim 12, wherein the growth hormone is administered in combination with a substance that leads to GH stimulation.
- 14. (New) The method of claim 12, wherein the growth hormone is administered in combination with a substance that induces a GH analogous effect.
- 15. (New) The method of claim 12, wherein the growth hormone is administered in combination with a substance that promotes IGF-I release.
- 16. (New) A method of treating female and male sexual functional disorders comprising administering an effective amount of a substance that leads to growth hormone stimulation, wherein the sexual functional disorder is manifested by lack or loss of libido, problems relating to orgasms, insufficient lubrication, or erectile dysfunction and wherein insufficient increase in hGH concentration occurs during sexual stimulation or a hGH deficiency exists.
- 17. (New) A method of treating female and male sexual functional disorders comprising administering an effective amount of a substance that induces a growth hormone analogous effect, wherein the sexual functional disorder is manifested by lack or loss of libido, problems relating to orgasms, insufficient lubrication, or erectile dysfunction and wherein

insufficient increase in hGH concentration occurs during sexual stimulation or a hGH deficiency exists.

18. (New) A method of treating female and male sexual functional disorders comprising administering an effective amount of a substance that promotes IGF-1 release, wherein the sexual functional disorder is manifested by lack or loss of libido, problems relating to orgasms, insufficient lubrication, or erectile dysfunction and wherein insufficient increase in hGH concentration occurs during sexual stimulation or a hGH deficiency exists.

## REMARKS

The above preliminary amendment is made to have the specification and claims conform to United States Patent and Trademark Office practice. Please refer to the Marked-Up Version attached herewith.

If a telephone conference would be helpful in resolving any issues concerning this communication, please contact Applicants' primary attorney-of record, Ronald A. Daignault (Reg. No. 25,968), at (612) 371.5381.

Please charge any additional fees or credit any overpayment to Deposit Account No. 13-2725.

Respectfully submitted,

MERCHANT & GOULD P.C.

P.O. Box 2903

Minneapolis, Minnesota 55402-0903

(612) 332-5300

Dated: March 21, 2002

RAD/rw

Ronald A. Daignault Reg. No. 25,968

23552
PATENT TRADEMARK OFFICE

#### (MARKED UP VERSION)

#### Administration of Growth Hormone (hGH) for Therapy of Sexual Functional Disorders

#### **Technical Field**

[001] This invention concerns the introduction of human growth hormone (hGH, GH) for the manufacture of medicaments for the treatment of sexual functional disorders in both male and female patients as well as the specific methods of treatment undertaken.

#### **Background of the Invention**

[002] Symptoms indicating sexual functional disorders are, for example, lack or loss of libido, problems relating to orgasms, insufficient lubrication and erectile dysfunction (ED). It could be deduced from certain cases that the basic aetiologies leading to these sexual functional disorders were due to a number of different reasons. Apart from the cases of mixed aetiologies, vascular (arterial, venous), psychogenic, neurogenic, medicamentous-induced and cavernous sexual functional disorders are also differentiated.

[003] Taking into consideration the nature of the underlying aetiologies, causal therapy of the sexual functional disorders is undertaken whenever possible. Up till now this method of therapy has only proved successful in the rare cases (e.g. by psychotherapy, hormone treatment, a change-over of medication) so that the main methods of therapy still remain unspecific.

[004] There are many different methods of therapy [are] available for males with ED compared to those available for females with sexual functional disorders. These include oral, topical, intracavernous, intraurethral and also a combination of drugs. These methods do not constitute a causal therapy, rather the aim is to achieve a direct or indirect relaxation (flaccidity) of the corpus cavernosum smooth musculature and the penile arteries. Together with an increase of blood circulation, penile erection is achieved. Furthermore, the vacuum pump, arterial shunt procedure, venous closure operations and penile prosthesis implantations are also methods used for therapy. Until the introduction of sildenafil (VIAGRA®), the most widely used form of therapy involved the administration of intracavernous vasoactive substances. At the present time, sildenafil is used as the so-called "first line therapy" providing there are no known contraindications. The oral phosphodiesterase type 5 inhibitor (PDE5) does not provide a basis for causal therapy. With the inhibition of PDE5, hydrolysis

of cyclic guanosinmonophosphate (cGMP) of an intracellular second messenger is prohibited, resulting in relaxation of the corpus cavernosum smooth musculature. This effective mechanism is assumed beneficial for the increase of lubrication in women, however, recent studies have yet to prove its effectivity.

## **Summary of the Invention**

[005] The aim of this innovative breakthrough was to present a new therapy for both males and females suffering from sexual functional disorders.

[006] Surprisingly, it has been shown that the growth hormone (hGH) plays an essential role in sexual stimulation, as an enormous unexpected increase of this hormone was seen to be present at the onset of sexual stimulation.

[007] The focal point of this breakthrough is therefore the use of hGH for the manufacture of medicaments for the treatment of sexual functional disorders in both males and females with e.g. lack or loss of libido, problems relating to orgasms, insufficient lubrication and erectile dysfunction and for the therapy of the aforesaid functional disorders.

[008] Accordingly, the present invention is a method of treating female and male sexual functional disorders that includes administering an effective amount of growth hormone. The sexual functional disorder is manifested by a lack or loss of libido, problems relating to orgasms, insufficient lubrication, or erectile dysfunction; there is an insufficient increase in hGH concentration occurs during sexual stimulation or a hGH deficiency exists. A further [issue] aspect is the use of hGH for the therapy of functional disorders in synergic combination with effective substances which result in GH stimulation, induce a GH analogous effect or promote IGF-I release.

[009] [The scientific results described are emphasized as follows:]

## **Brief Description of the Drawings**

[010] <u>Figure 1</u> shows the average values and standard deviations of the growth hormone (hGH) concentrations (ng/ml) in cavernous and peripheral blood samples taken from 35 healthy probands during the four different phases of the penile erectile tissue (flaccidity, tumescence, rigidity and detumescence).

[011] <u>Figure 2</u> shows the average values and standard deviations of the growth hormone (hGH) concentrations (ng/ml) in the cavernous and peripheral blood samples during the three different penile phases (flaccidity, tumescence and detumescence) in 36 patients with erectile dysfunction. Rigidity was not achieved due to disorder of patient. The axis scale was selected as in Figure 1 in order to demonstrate clearly the difference (P<0.05).

[012] <u>Figure 3</u> shows the average values and standard deviations of the dose-dependent decrease in relaxation of 12 human corpus cavernosum strips after application of recombinant hGH.

[013] <u>Figure 4</u> shows the average values and standard deviations of dose-dependent increase of cyclic guanosinmonophasphate (cGMP) of 3 human corpus cavernosum strips respectively, following incubation with recombinant hGH or sodium nitroprusside (SNP). Incubation with SNP was carried out using a concentration of 0.01 and 1  $\mu$ Mol. For this reason, no value for SNP was achieved with 0.0001  $\mu$ Mol.

## **Detailed Description of the Invention**

[014] In order to gain a better understanding of the physiology of a naturally induced erection and the pathophysiology of erectile dysfunction, a new method of investigation was developed. This involved detection of endogenous human neurotransmitters, neuromodulators and hormones which might have some connection with an erection or its sexual function. These new methods of investigation aim to improve the diagnostics and therapy (application of endogenous substances and causal therapy) for those patients with sexual functional disorders.

[015] Blood was taken simultaneously from the corpus cavernosum (CC, cavernous) and the cubital vein (CV, peripheral) in 35 healthy probands during the phases of flaccidity, tumescence, rigidity and detumescence. Audiovisual and tactile means were then provided to aid sexual stimulation (Figure 1). The procedure involving 36 patients with ED was identical to that of the healthy probands with the exception of blood withdrawal during the rigidity phase, (this penile erection phase cannot be achieved in patients with ED) (Figure 2). The hGH concentrations were determined with an immunoradiometric assay (IRMA). This form of investigation resulted in several new findings.

[016] [1.]The highest increase of hGH concentration was found during tumescence, namely the point in time when sexual stimulation is at its peak. [2.]The peripheral and cavernous hGH concentrations showed no significant differences in direct comparison with all penile phases. Peripheral blood withdrawal proved sufficient. [3.]When comparing the healthy probands with the patients, there were significant differences with regards to the hGH concentrations, in particular, a significantly reduced increase of the hGH concentration during the tumescence phase.

[017] These data show for the first time the surprising causal connection of hGH formed by the hypophysis with sexual stimulation and the resulting penile erection. The reduced expression of hGH on sexual stimulation in patients is further proof for the significance of this hormone, the lack of which is connected with sexual functional disorders and erectile dysfunction in this study.

[018] By means of extensive in vitro investigations using human corpus cavernosum (CC) tissue as well as the in vivo results described, important indications could be deduced for the possible physiological connections between hGH and penile erection. [1.]Organ bath experiments (in vitro method to evaluate the relaxing properties of substances) with human CC were carried out to assess dose-dependent relaxations following application of hGH (Figure 3).

[019] [2.]Incubation experiments (in vitro method to evaluate the content of cyclic nucleotides in tissues in response to drug exposition, in this case cGMP, after incubation with various substances) with human CC were shown to have dose-dependent higher cGMP concentrations after application of hGH than was the case after incubation with sodium nitroprusside (SNP), a classic NO donator (Figure 4).

[020] Based on our human findings, it can be assumed that hGH plays a decisive role in sexual function (sexual stimulation), in particular, in penile erection. Furthermore, it was shown that the peripheral reaction of hGH induced an increase of cGMP, thereby physiologically forming a link between relaxation of CC with that of the ensuing erection. Due to the anatomical similarities in the structure of the penis and the clitoris and the physiological conformities regarding sexual stimulation (e.g. congestion of the genital organs mediated by neurotransmitter on relaxation of the smooth musculature), the described reaction

of hGH in males must also apply to females too, since hGH is produced by the hypophysis in both sexes, therefore the same effect must also be evident in both sexes.

[021] With reference to the effect of hGH, it is already known that hGH does not focus on any particular tissue and that the activity and metabolism (anabolic) is increased in different tissues in both men and women. The growth hormone stimulates, for example, body growth (substitution in insufficient hGH-caused hyposomia) and protein metabolism (possible indication in cachexia, severe burn injuries and also anabolic abuse). Under the influence of hGH, an insulin-like growth factor I (IGF-I) is formed mainly in the liver but also in other tissues. This polypeptide (IGF-I) plays a significant mediating role in the process induced by hGH (Merimee T.J. and Grant M.B.: Growth hormone and its disorders. In: Principles and Practice of Endocrinology and Metabolism. Edited by Becker, K.L., Philadelphia, J.B. Lippincott Company, pp. 125-134, 1990).

[022] The most recent findings in humans show that there is a systemic increase of NO (nitric oxide) and cGMP under a substitution of recombinant produced hGH (r-hGH) in patients with hGH deficiency (Böger R.H. et al.: Nitric oxide may mediate the hemodynamic effects of recombinant growth hormone in patients with acquired growth hormone deficiency. J. Clin. Invest. 98: 2706-2713, 1996). This NO-cGMP path presents a very important positive significance in achieving penile erection (Burnett A.L. et al.: Nitric oxide: a physiologic mediator or penile erection. Science 257: 905, 1993). Likewise, recent findings derived from animal experiments using rats were able to show that an increase of NOS (nitric oxide synthase)-containing nerves (generating NO) in CC and dorsal penile nerves occurred under substitution of hGH. This took place despite initiation of neurogenic damage some weeks before (Jung G.W. et al.: Growth hormone enhances regeneration of nitric oxide synthasecontaining penile nerves after cavernous neurotomy in rats. J. Urol. 160: 1899-1904, 1998). The patent WO98/42361 (Human erectile dysfunction and methods of treatment) stems from these results and describes the indication of hGH therapy for the prevention and treatment of neurogenic erectile dysfunction of different aetiologies (condition following extensive pelvic operations or pelvic trauma, diabetes, alcoholism and aging process).

[023] Our findings on human models show for the first time a positive causal connection between sexual stimulation, increase of hGH and penile erection. The reduced (often totally lacking) increase of hGH in patients with ED emphasizes the importance of this hormone.

The in vitro data conclude that the NO-cGMP path is activated by hGH, leading to relaxation of the CC, thus resulting in penile erection.

[024] The appropriate therapy for all patients (both sexes) with sexual functional disorders includes peripheral blood sample to determine the basal hGH concentration. This form of therapy is undertaken independent of the underlying aetiology(ies). Following this, a further blood sample is taken under sexual stimulation (audiovisual, tactile) in order to detect the stimulated hGH concentration. In cases of insufficient or no reaction at all to sexual stimulation (e.g. lubrication, penile erection) and inadequate increase of hGH concentration, a continuous, strictly controlled therapy with hGH should ensue for a certain period of time (e.g. 2 - 6 months).

[025] Suitable pharmaceutical preparations for therapy include solid or liquid forms of administration for oral intake, such as tablets, capsules or emulsions, parenteral forms of administration for injection or non-invasive application or transdermal topical systems, such as plasters, creams, gels, lotions or transdermal films. The administered amount for successful therapy lies between 0.01 and 500 mg per dosage unit; recommended is between 0.1 and 100 mg.

[026] Improvement of therapy outcome can be achieved by administration of a combination of medicaments containing, besides hGH, a synergic combination of substances which lead to GH stimulation, induce a GH analogous effect or promote IGF-I release.

[027] These substances do not have to be combined into one particular medication, but can be administered in separate suitable galenic preparations to be taken at the same time or taken separately according to the specific course of therapy. It is essential that the specialist instructs and informs the patient with regards to the suitable dosage or in which combination the medicaments should be taken, likewise which substance should be administered to ensure the best possible therapy outcome. Furthermore, it is permissible to combine several of the named substances to treat the individual patient accordingly.

[028] The suitable substances to be used as a combination therapy in order to achieve GH stimulation are familiar to the specialist. For example, arginine, alpha 1 and alpha 2-agonists, such as clonidine, norepinephrine or salbutamol, glucagon, pyridostigmine, galanine, GH-

releasing hormone, NPY (neuropeptide Y) and dopamine agonists, such as apomorphine, quinpirole or cabergoline.

[029] Suitable substances which induce a GH analogous effect include, for instance, GHRP (growth hormone, releasing hexapeptide, hexareline), GH releasing peptide 1, 2, 6 and non-peptidergic agonists of growth hormone releasing peptide such as MK 0677, EP 51389 (2-methylalanyl-2-methyl-D-tryptophyl-2-methyl-D-tryptophanamide), L 692429 (3-amino-3-methyl-N-[(3R)-2,3,4,5-tetrahydro-2-oxo-1-[[1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-1H-1-benzazepine-3-yl]butanamide), or L 692585 (3-[[(2R)-2hydroxypropyl]amino]-3-methyl-N-[(3R)-2,3,4,5-tetrahydro-2-oxo-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-1H-1-benzazepine-3-yl]-butanamide).

[030] Suitable substances which promote IGF-I release include, for example, cannabinoide such as e.g. HU-210 (3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-6H-dibenzo[b,d]pyran-9-methanol) or serotonine receptor agonists such as e.g. 8-OH DPAT (8-hydroxy-2-dipropylamino)tetraline), or SC 53116 (4-amino-5-chloro-N-[[(1s,7aS)-hexahydro-1H-pyrrolizine-1-yl]methyl]-2-methoxy-benzamide).

#### WE CLAIM:

Please cancel claims 1-11 without prejudice or disclaimer and add claims 12-18.

- 12. (New) A method of treating femaled and male sexual functional disorders comprising administering an effective amount of a growth hormone, wherein the sexual functional disorder is manifested by lack or loss of libido, problems relating to orgasms, insufficient lubrication, or erectile dysfunction and wherein insufficient increase in hGH concentration occurs during sexual stimulation or a hGH deficiency exists.
- 13. (New) The method of claim 12, wherein the growth hormone is administered in combination with a substance that leads to GH stimulation.
- 14. (New) The method of claim 12, wherein the growth hormone is administered in combination with a substance that induces a GH analogous effect.
- 15. (New) The method of claim 12, wherein the growth hormone is administered in combination with a substance that promotes IGF-I release.
- 16. (New) A method of treating female and male sexual functional disorders comprising administering an effective amount of a substance that leads to growth hormone stimulation, wherein the sexual functional disorder is manifested by lack or loss of libido, problems relating to orgasms, insufficient lubrication, or erectile dysfunction and wherein insufficient increase in hGH concentration occurs during sexual stimulation or a hGH deficiency exists.
- 17. (New) A method of treating female and male sexual functional disorders comprising administering an effective amount of a substance that induces a growth hormone analogous effect, wherein the sexual functional disorder is manifested by lack or loss of libido, problems relating to orgasms, insufficient lubrication, or erectile dysfunction and wherein insufficient increase in hGH concentration occurs during sexual stimulation or a hGH deficiency exists.
- 18. (New) A method of treating female and male sexual functional disorders comprising administering an effective amount of a substance that promotes IGF-1 release,

wherein the sexual functional disorder is manifested by lack or loss of libido, problems relating to orgasms, insufficient lubrication, or erectile dysfunction and wherein insufficient increase in hGH concentration occurs during sexual stimulation or a hGH deficiency exists.

Rec'd PCT/PTO 21 MAR 2002 10/018719

Rambergstr. 29 D-30161 Hannover Tel.:+49 511 3 18 04 73

14.12.01

# To whom it may concern

I, Nicola van Domick, Personal Assistant, with residence at Rambergstr. 29, 30161 Hannover, Germany, herewith declare that I am well acquainted with the German and the English language and that the translated document is a true, correct, and complete English translation of the German International Patent Application No. WO 00/78328.

Hannover, Germany, 14.12.01

Nicola van Dornick

Rec'd PCT/PTO 21 MAR 2002 10/018719 #3

# Administration of Growth Hormone (hGH) for Therapy of Sexual Functional Disorders

This invention concerns the introduction of human growth hormone (hGH, GH) for the manufacture of medicaments for the treatment of sexual functional disorders in both male and female patients as well as the specific methods of treatment undertaken.

Symptoms indicating sexual functional disorders are, for example, lack or loss of libido, problems relating to orgasms, insufficient lubrication and erectile dysfunction (ED). It could be deduced from certain cases that the basic aetiologies leading to these sexual functional disorders were due to a number of different reasons. Apart from the cases of mixed aetiologies, vascular (arterial, venous), psychogenic, neurogenic, medicamentous-induced and cavernous sexual functional disorders are also differentiated.

Taking into consideration the nature of the underlying aetiologies, causal therapy of the sexual functional disorders is undertaken whenever possible. Up till now this method of therapy has only proved successful in the rare cases (e.g. by psychotherapy, hormone treatment, a change-over of medication) so that the main methods of therapy still remain unspecific.

There are many different methods of therapy are available for males with ED compared to those available for females with sexual functional disorders. These include oral, topical, intracavernous, intraurethral and also a combination of drugs. These methods do not constitute a causal therapy, rather the aim is to achieve a direct or indirect relaxation (flaccidity) of the corpus cavernosum smooth musculature and the penile arteries. Together with an increase of blood circulation, penile erection is achieved. Furthermore, the vacuum pump, arterial shunt procedure, venous closure operations and penile prosthesis implantations are also methods used for therapy. Until the introduction of sildenafil (VIAGRA®), the most widely used form of therapy involved the administration of intracavernous vasoactive substances. At the present time, sildenafil is used as the so-called "first line therapy" providing there are no known contraindications. The oral phosphodiesterase type 5 inhibitor (PDE5) does not provide a basis for causal therapy. With the inhibition of PDE5, hydrolysis of cyclic guanosinmonophosphate (cGMP) of an intracellular second messenger is prohibited, resulting in relaxation of the corpus cavernosum smooth musculature. This effective mechanism is assumed beneficial for the increase of lubrication in women, however, recent studies have yet to prove its effectivity.

The aim of this innovative breakthrough was to present a new therapy for both males and females suffering from sexual functional disorders.

Surprisingly, it has been shown that the growth hormone (hGH) plays an essential role in sexual stimulation, as an enormous unexpected increase of this hormone was seen to be present at the onset of sexual stimulation.

The focal point of this breakthrough is therefore the use of hGH for the manufacture of medicaments for the treatment of sexual functional disorders in both males and females with e.g. lack or loss of libido, problems relating to orgasms, insufficient lubrication and erectile dysfunction and for the therapy of the aforesaid functional disorders.

A further issue is the use of hGH for the therapy of functional disorders in synergic combination with effective substances which result in GH stimulation, induce a GH analogous effect or promote IGF-I release.

The scientific results described are emphasized as follows:

Figure 1 shows the average values and standard deviations of the growth hormone (hGH) concentrations (ng/ml) in cavernous and peripheral blood samples taken from 35 healthy probands during the four different phases of the penile erectile tissue (flaccidity, tumescence, rigidity and detumescence).

Figure 2 shows the average values and standard deviations of the growth hormone (hGH) concentrations (ng/ml) in the cavernous and peripheral blood samples during the three different penile phases (flaccidity, tumescence and detumescence) in 36 patients with erectile dysfunction. Rigidity was not achieved due to disorder of patient. The axis scale was selected as in Figure 1 in order to demonstrate clearly the difference (P<0.05).

Figure 3 shows the average values and standard deviations of the dose-dependent decrease in relaxation of 12 human corpus cavernosum strips after application of recombinant hGH.

Figure 4 shows the average values and standard deviations of dose-dependent increase of cyclic guanosinmonophasphate (cGMP) of 3 human corpus cavernosum strips respectively, following incubation with recombinant hGH or sodium nitroprusside (SNP). Incubation with SNP was carried out using a concentration of 0.01 and 1  $\mu$ Mol. For this reason, no value for SNP was achieved with 0.0001  $\mu$ Mol.

In order to gain a better understanding of the physiology of a naturally induced erection and the pathophysiology of erectile dysfunction, a new method of investigation was developed. This involved detection of endogenous human neurotransmitters, neuromodulators and hormones which might have some connection with an erection or its sexual function. These

new methods of investigation aim to improve the diagnostics and therapy (application of endogenous substances and causal therapy) for those patients with sexual functional disorders.

Blood was taken simultaneously from the corpus cavernosum (cavernous) and the cubital vein (peripheral) in 35 healthy probands during the phases of flaccidity, tumescence, rigidity and detumescence. Audiovisual and tactile means were then provided to aid sexual stimulation (Figure 1). The procedure involving 36 patients with ED was identical to that of the healthy probands with the exception of blood withdrawal during the rigidity phase, (this penile erection phase cannot be achieved in patients with ED) (Figure 2). The hGH concentrations were determined with an immunoradiometric assay (IRMA). This form of investigation resulted in several new findings.

- 1. The highest increase of hGH concentration was found during tumescence, namely the point in time when sexual stimulation is at its peak.
- 2. The peripheral and cavernous hGH concentrations showed no significant differences in direct comparison with all penile phases. Peripheral blood withdrawal proved sufficient.
- 3. When comparing the healthy probands with the patients, there were significant differences with regards to the hGH concentrations, in particular, a significantly reduced increase of the hGH concentration during the tumescence phase.

These data show for the first time the surprising causal connection of hGH formed by the hypophysis with sexual stimulation and the resulting penile erection. The reduced expression of hGH on sexual stimulation in patients is further proof for the significance of this hormone, the lack of which is connected with sexual functional disorders and erectile dysfunction in this study.

By means of extensive in vitro investigations using human corpus cavernosum (CC) tissue as well as the in vivo results described, important indications could be deduced for the possible physiological connections between hGH and penile erection.

- Organ bath experiments (in vitro method to evaluate the relaxing properties of substances) with human CC were carried out to assess dose-dependent relaxations following application of hGH (Figure 3).
- 2. Incubation experiments (in vitro method to evaluate the content of cyclic nucleotides in tissues in response to drug exposition, in this case cGMP, after incubation with various

substances) with human CC were shown to have dose-dependent higher cGMP concentrations after application of hGH than was the case after incubation with sodium nitroprusside (SNP), a classic NO donator (Figure 4).

Based on our human findings, it can be assumed that hGH plays a decisive role in sexual function (sexual stimulation), in particular, in penile erection. Furthermore, it was shown that the peripheral reaction of hGH induced an increase of cGMP, thereby physiologically forming a link between relaxation of CC with that of the ensuing erection. Due to the anatomical similarities in the structure of the penis and the clitoris and the physiological conformities regarding sexual stimulation (e.g. congestion of the genital organs mediated by neurotransmitter on relaxation of the smooth musculature), the described reaction of hGH in males must also apply to females too, since hGH is produced by the hypophysis in both sexes, therefore the same effect must also be evident in both sexes.

With reference to the effect of hGH, it is already known that hGH does not focus on any particular tissue and that the activity and metabolism (anabolic) is increased in different tissues in both men and women. The growth hormone stimulates, for example, body growth (substitution in insufficient hGH-caused hyposomia) and protein metabolism (possible indication in cachexia, severe burn injuries and also anabolic abuse). Under the influence of hGH, an insulin-like growth factor I (IGF-I) is formed mainly in the liver but also in other tissues. This polypeptide (IGF-I) plays a significant mediating role in the process induced by hGH (Merimee T.J. and Grant M.B.: Growth hormone and its disorders. In: Principles and Practice of Endocrinology and Metabolism. Edited by Becker, K.L., Philadelphia, J.B. Lippincott Company, pp. 125-134, 1990).

The most recent findings in humans show that there is a systemic increase of NO (nitric oxide) and cGMP under a substitution of recombinant produced hGH (r-hGH) in patients with hGH deficiency (Böger R.H. et al.: Nitric oxide may mediate the hemodynamic effects of recombinant growth hormone in patients with acquired growth hormone deficiency. J. Clin. Invest. 98: 2706-2713, 1996). This NO-cGMP path presents a very important positive significance in achieving penile erection (Burnett A.L. et al.: Nitric oxide: a physiologic mediator or penile erection. Science 257: 905, 1993). Likewise, recent findings derived from animal experiments using rats were able to show that an increase of NOS (nitric oxide synthase)-containing nerves (generating NO) in CC and dorsal penile nerves occurred under substitution of hGH. This took place despite initiation of neurogenic damage some weeks before (Jung G.W. et al.: Growth hormone enhances regeneration of nitric oxide synthase-containing penile nerves after cavernous neurotomy in rats. J. Urol. 160: 1899-1904, 1998). The patent WO98/42361 (Human erectile dysfunction and methods of treatment) stems from these results and describes the indication of hGH therapy for the prevention and treatment of

neurogenic erectile dysfunction of different aetiologies (condition following extensive pelvic operations or pelvic trauma, diabetes, alcoholism and aging process).

Our findings on human models show for the first time a positive causal connection between sexual stimulation, increase of hGH and penile erection. The reduced (often totally lacking) increase of hGH in patients with ED emphasizes the importance of this hormone. The in vitro data conclude that the NO-cGMP path is activated by hGH, leading to relaxation of the CC, thus resulting in penile erection.

The appropriate therapy for all patients (both sexes) with sexual functional disorders includes peripheral blood sample to determine the basal hGH concentration. This form of therapy is undertaken independent of the underlying aetiology(ies). Following this, a further blood sample is taken under sexual stimulation (audiovisual, tactile) in order to detect the stimulated hGH concentration. In cases of insufficient or no reaction at all to sexual stimulation (e.g. lubrication, penile erection) and inadequate increase of hGH concentration, a continuous, strictly controlled therapy with hGH should ensue for a certain period of time (e.g. 2 - 6 months).

Suitable pharmaceutical preparations for therapy include solid or liquid forms of administration for oral intake, such as tablets, capsules or emulsions, parenteral forms of administration for injection or non-invasive application or transdermal topical systems, such as plasters, creams, gels, lotions or transdermal films. The administered amount for successful therapy lies between 0.01 and 500 mg per dosage unit; recommended is between 0.1 and 100 mg.

Improvement of therapy outcome can be achieved by administration of a combination of medicaments containing, besides hGH, a synergic combination of substances which lead to GH stimulation, induce a GH analogous effect or promote IGF-I release.

These substances do not have to be combined into one particular medication, but can be administered in separate suitable galenic preparations to be taken at the same time or taken separately according to the specific course of therapy. It is essential that the specialist instructs and informs the patient with regards to the suitable dosage or in which combination the medicaments should be taken, likewise which substance should be administered to ensure the best possible therapy outcome. Furthermore, it is permissible to combine several of the named substances to treat the individual patient accordingly.

The suitable substances to be used as a combination therapy in order to achieve GH stimulation are familiar to the specialist. For example, arginine, alpha 1 and alpha 2-agonists,

such as clonidine, norepinephrine or salbutamol, glucagon, pyridostigmine, galanine, GH-releasing hormone, NPY (neuropeptide Y) and dopamine agonists, such as apomorphine, quinpirole or cabergoline.

Suitable substances which induce a GH analogous effect include, for instance, GHRP (growth hormone, releasing hexapeptide, hexareline), GH releasing peptide 1, 2, 6 and non-peptidergic agonists of growth hormone releasing peptide such as MK 0677, EP 51389 (2-methylalanyl-2-methyl-D-tryptophyl-2-methyl-D-tryptophanamide), L 692429 (3-amino-3-methyl-N-[(3R)-2,3,4,5-tetrahydro-2-oxo-1-[[1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-1H-1-benzazepine-3-yl]butanamide], or L 692585 (3-[[(2R)-2hydroxypropyl]amino]-3-methyl-N-[(3R)-2,3,4,5-tetrahydro-2-oxo-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-1H-1-benzazepine-3-yl]-butanamide).

Suitable substances which promote IGF-I release include, for example, cannabinoide such as e.g. HU-210 (3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-6H-dibenzo[b,d]pyran-9-methanol) or serotonine receptor agonists such as e.g. 8-OH DPAT (8-hydroxy-2-dipropylamino)tetraline), or SC 53116 (4-amino-5-chloro-N-[[(1s,7aS)-hexahydro-1H-pyrrolizine-1-yl]methyl]-2-methoxy-benzamide).

#### **Patent Claims**

- Use of growth hormone for manufacture of medicaments for the therapy of female and male sexual functional disorders, such as the lack or loss of libido, problems relating to orgasms, insufficient lubrication, erectile dysfunction for all aetiologies not treated in a causal manner, in cases of insufficient increase in hGH concentration during sexual stimulation and for those with hGH deficiency.
- 2. Use of growth hormone according to Claim 1 in combination with substances which lead to GH stimulation.
- 3. Use of growth hormone according to Claim 1 in combination with substances which induce a GH analogous effect.
- 4. Use of growth hormone according to Claim 1 in combination with substances which promote IGF-I release.
- 5. Use of substances which lead to GH stimulation according to Claim 1.
- 6. Use of substances which induce a GH analogous effect according to Claim 1.
- 7. Use of substances which promote IGF-I release according to Claim 1.
- 8. Use of growth hormone for the therapy of female and male sexual functional disorders such as lack or loss of libido, problems relating to orgasms, insufficient lubrication, erectile dysfunction for all aetiologies not treated in a causal manner, in cases of insufficient increase in hGH concentration during sexual stimulation and for those with hGH deficiency.
- 9. Use of substances which lead to GH stimulation according to Claim 8.
- 10. Use of substances which induce a GH analogous effect according to Claim 8.
- 11. Use of substances which promote IGF-I release according to Claim 8.

## **Summary**

This innovative breakthrough concerns the use of human growth hormone (hGH, GH) alone or in combination with substances which result in GH stimulation, induce a GH analogous effect or promote IGF-I release for the manufacture of medicaments for the treatment of sexual functional disorders of both sexes and the methods of therapy of the named disorders.

Figure 1: Course of human growth hormone levels in the cavernous (CC = Corpus cavernosum) and the systemic blood (CV = cubital vein) of healthy males during penile flaccidity, tumescence, rigidity, and detumescence. All data are given in ng/mL as mean  $\pm$  standard deviation of the mean.

**Figure 2:** Course of hGH serum levels in blood samples taken during different penile conditions (flaccidity, tumescence, detumescence) from the corpus cavernosum (CC) and the cubital vein (CV) of patients suffering from erectile dysfunction.

Figure 3: Relaxation of human corpus cavernosum strips in vitro induced by cumulative addition of recombinant human growth hormone. Each point is expressed as percentage of maximum norepinephrine-HCl - induced tension and represents mean  $\pm$  standard deviation of the mean of n = 12 determinations.

**Figure 4:** Stimulating effects of recombinant human growth hormone (rhGH) and sodiumnitroprusside (SNP) on tissue levels of cGMP in isolated human cavernous smooth muscle strips. Each bar represents mean ± standard deviation of the mean of n = 3 - 6 determinations.

Inventor: Becker, et al.
Docket No· 13390.2USWO
Title: USE OF GROWTH HORMONE (HGH) FOR THE TREATMENT OF SEXUAL FUNCTIONAL DISTURBANCES
Senal No· 10/018719
Sheet 1 of 2

Figure 1





Figure 2



Inventor: Becker, et al.
Docket No.: 13390.2USWO
Title: USE OF GROWTH HORMONE (HGH) FOR THE TREATMENT OF SEXUAL
FUNCTIONAL DISTURBANCES
Serial No.: 10/018719
Sheet 2 of 2

Figure 3





Figure 4



#### MERCHANT & GOULD P.C.

# **United States Patent Application**

## COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor I hereby declare that: my residence, post office address and citizenship are as stated below next to my name; that

I verily believe I am the original, first and sole inventor (if only one name is listed below) or a joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: USE OF GROWTH HORMONE (HGH) FOR THE TREATMENT OF SEXUAL FUNCTIONAL DISTURBANCES

The specification of which			
a. is attached hereto b. was filed on as application described and claimed in internation which I have reviewed and for which	on serial no. and was amen nal no. PCT/EP00/05517 filed ch I solicit a United States pate	June 15, 2000 and as a	e) (in the case of a PCT-filed application) mended on (if any),
I hereby state that I have reviewed any amendment referred to above.	and understand the contents of	the above-identified sp	pecification, including the claims, as amended by
that of the application on the basis a no such applications have be such applications have been	so identified below any foreign of which priority is claimed: en filed. filed as follows:	application for patent	any foreign application(s) for patent or inventor's or inventor's certificate having a filing date before
	IGN APPLICATION(S), IF ANY, (	CLAIMING PRIORITY U	NDER 35 USC § 119
<u>CO</u> UNTRY	APPLICATION NUMBER	DATE OF FILING	DATE OF ISSUE
GERMANY	199 27 678.1	(day, month, year)	(day, month, year)
		17/JUNE/1	
COUNTRY	GN APPLICATION(S), IF ANY, F		ORITY APPLICATION(S)
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)
manner provided by the first paragram	atter of each of the claims of the aph of Title 35, United States ( Regulations, § 1.56(a) which o	is application is not dis Code, § 112. I acknowle	rates and PCT international application(s) listed aclosed in the prior United States application in the edge the duty to disclose material information as iling date of the prior application and the national
U.S. APPLICATION NUMBER	DATE OF FILING	(day, month, year)	STATUS (patented, pending, abandoned)
I hereby claim the benefit under Titl	e 35, United States Code § 119	θ(e) of any United State	es provisional application(s) listed below:
U.S. PROVISIONAL API			TE OF FILING (Day, Month, Year)

I acknowledge the duty to disclose information that is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56 (reprinted below):

## § 1.56 Duty to disclose information material to patentability.

n.

- (a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is canceled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is canceled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced or attempted or the duty of disclosure was violated through bad faith or intentional misconduct. The Office encourages applicants to carefully examine:
  - (1) prior art cited in search reports of a foreign patent office in a counterpart application, and
- (2) the closest information over which individuals associated with the filing or prosecution of a patent application believe any pending claim patentably defines, to make sure that any material information contained therein is disclosed to the Office.
- (b) Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and
  - (1) It establishes, by itself or in combination with other information, a prima facie case of unpatentability of a claim;
  - (2) It refutes, or is inconsistent with, a position the applicant takes in:
    - (i) Opposing an argument of unpatentability relied on by the Office, or
    - (ii) Asserting an argument of patentability.

A prima facie case of unpatentability is established when the information compels a conclusion that a claim is unpatentable under the preponderance of evidence, burden-of-proof standard, giving each term in the claim its broadest reasonable construction consistent with the specification, and before any consideration is given to evidence which may be submitted in an attempt to establish a contrary conclusion of patentability.

- (c) Individuals associated with the filing or prosecution of a patent application within the meaning of this section are:
  - (1) Each inventor named in the application:
  - (2) Each attorney or agent who prepares or prosecutes the application; and
- (3) Every other person who is substantively involved in the preparation or prosecution of the application and who is associated with the inventor, with the assignee or with anyone to whom there is an obligation to assign the application.
- (d) Individuals other than the attorney, agent or inventor may comply with this section by disclosing information to the attorney, agent, or inventor.
- (e) In any continuation-in-part application, the duty under this section includes the duty to disclose to the Office all information known to the person to be material to patentability, as defined in paragraph (b) of this section, which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby appoint the following attorney(s) and/or patent agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected herewith:

Albrecht, John W.	Reg. No. 40,481	Leonard, Christopher J.	Reg. No. 41,940
Ali, M. Jeffer	Reg. No. 46,359	Liepa, Mara E.	Reg. No. 40,066
Altera, Allan G.	Reg. No. 40,274	Lindquist, Timothy A.	Reg. No. 40,701
Anderson, Gregg I.	Reg. No. 28,828	Lown, Jean A.	Reg. No. 48,428
Batzli, Brian H.	Reg. No. 32,960	Mayfield, Denise L.	Reg. No. 33,732
Beard, John L.	Reg. No. 27,612	McDonald, Daniel W.	Reg. No. 32,044
Berns, John M.	Reg. No. 43,496	McIntyre, Jr., William F.	Reg. No. 44,921
Branch, John W.	Reg. No. 41,633	Mitchem, M. Todd	Reg. No. 40,731
Brown, Jeffrey C.	Reg. No. 41,643	Mueller, Douglas P.	Reg. No. 30,300
Bruess, Steven C.	Reg. No. 34,130	Nelson, Anna M.	Reg. No. 48,935
Byrne, Linda M.	Reg. No. 32,404	Paley, Kenneth B.	Reg. No. 38,989
Campbell, Keith	Reg. No. 46,597	Parsons, Nancy J.	Reg. No. 40,364
Carlson, Alan G.	Reg. No. 25,959	Pauly, Daniel M.	Reg. No. 40,123
Caspers, Philip P.	Reg. No. 33,227	Phillips, John B.	Reg. No. 37,206
Clifford, John A.	Reg. No. 30,247	Pino, Mark J.	Reg. No. 43,858
Cook, Jeffrey	Reg. No. 48,649	Prendergast, Paul	Reg. No. 46,068
Daignault, Ronald A.	Reg. No. 25,968	Pytel, Melissa J.	Reg. No. 41,512
Daley, Dennis R.	Reg. No. 34,994	Qualey, Terry	Reg. No. 25,148
Daulton, Julie R.	Reg. No. 36,414	Reich, John C.	Reg. No. 37,703
DeVries Smith, Katherine M.	Reg. No. 42,157	Reiland, Earl D.	Reg. No. 25,767
DiPietro, Mark J.	Reg. No. 28,707	Samuels, Lisa A.	Reg. No. 43,080
Doscotch, Matthew A.	Reg No. P-48,957	Schmaltz, David G.	Reg. No. 39,828
Edell, Robert T.	Reg. No. 20,187	Schuman, Mark D.	Reg. No. 31,197
Epp Ryan, Sandra	Reg. No. 39,667	Schumann, Michael D.	Reg. No. 30,422
Glance, Robert J.	Reg. No. 40,620	Scull, Timothy B.	Reg. No. 42,137
Goff, Jared S.	Reg. No. 44,716	Sebald, Gregory A.	Reg. No. 33,280
Goggin, Matthew J.	Reg. No. 44,125	Skoog, Mark T.	Reg. No. 40,178
Goffa, Charles E.	Reg. No. 26,896	Spellman, Steven J.	Reg. No. 45,124
Gorman, Alan G.	Reg. No. 38,472	Stewart, Alan R.	Reg. No. 47,974
Gould, John D.	Reg. No. 18,223	Stoll-DeBell, Kirstin L.	Reg. No. 43,164
Gregson, Richard	Reg. No. 41,804	Sullivan, Timothy	Reg. No. 47,981
Gresens, John J.	Reg. No. 33,112	Sumner, John P.	Reg. No. 29,114
Hamer, Samuel A.	Reg. No. 46,754	Swenson, Erik G.	Reg. No. 45,147
Hamre, Curtis B.	Reg. No. 29,165	Tellekson, David K.	Reg. No. 32,314
Harrison, Kevin C.	Reg. No. 46,759	Trembath, Jon R.	Reg. No. 38,344
Heitzberg, Brett A.	Reg. No. 42,660	Tunheim, Marcia A.	Reg. No. 42,189
Hillson, Randall A.	Reg. No. 31,838	Underhill, Albert L.	Reg. No. 27,403
Holzer, Jr., Richard J.	Reg. No. 42,668	Vandenburgh, J. Derek	- · · · · · · · · · · · · · · · · · · ·
in the second se		•	Reg. No. 32,179
Hope, Leonard J.	Reg. No. 44,774	Wahl, John R.	Reg. No. 33,044
Jardine, John S.	Reg. No. P-48,835	Weaver, Paul L.	Reg. No. 48,640
Johns, Nicholas P.	Reg. No. 48,995	Welter, Paul A.	Reg. No. 20,890
Johnston, Scott W.	Reg. No. 39,721	Whipps, Brian	Reg. No. 43,261
Kadievitch, Natalie D.	Reg. No. 34,196	Whitaker, John E.	Reg. No. 42,222
Kaseburg, Frederick A.	Reg. No. 47,695	Wier, David D.	Reg. No. P-48,229
Kettelberger, Denise	Reg. No. 33,924	Williams, Douglas J.	Reg. No. 27,054
Keys, Jeramie J.	Reg. No. 42,724	Withers, James D.	Reg. No. 40,376
Knearl, Homer L.	Reg. No. 21,197	Witt, Jonelle	Reg. No. 41,980
Kowalchyk, Alan W.	Reg. No. 31,535	Wong, Thomas S.	Reg. No. 48,577
Kowalchyk, Katherine M.	Reg. No. 36,848	Wu, Tong	Reg. No. 43,361
Lacy, Paul E.	Reg. No. 38,946	Young, Thomas	Reg. No. 25,796
Larson, James A.	Reg. No. 40,443	Zeuli, Anthony R.	Reg. No. 45,255
*:			

I hereby authorize them to act and rely on instructions from and communicate directly with the person/assignee/attorney/firm/ organization who/which first sends/sent this case to them and by whom/which I hereby declare that I have consented after full disclosure to be represented unless/until I instruct Merchant & Gould P.C. to the contrary.

I understand that the execution of this document, and the grant of a power of attorney, does not in itself establish an attorney-client relationship between the undersigned and the law firm Merchant & Gould P.C., or any of its attorneys.

Please direct all correspondence in this case to Merchant & Gould P.C. at the address indicated below:

Merchant & Gould P.C. P.O. Box 2903 Minneapolis, MN 55402-0903



I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

		Full Name	Family Name	First Given Name	Second Given Name
_	20	Of Inventor	BECKER	Armin	Johannes
	0 1 Signa	Residence & Citizenship Mailing Address ture of Inventor 20 Full Name Of Inventor	City Hannover Address Blumhardthof 9	State or Foreign Country GERMANY City Hannover Date:	Country of Citizenship GERMANY  State & Zip Code/Country D-30625 / GERMANY
,	∂ † † † † † † † † † † † † † † † † † † †	Residence City & Citizenship Hemmingen  Mailing Address		State or Foreign Country GERMANY City	Country of Citizenship GERMANY  State & Zip Code/Country
		Address	Rehmenbreiten 6	Hemmingen	D-30966 / GERMANY
	Signa	ture of Inventor 20	Ju (7,2002		
1	Full Name Family Name Of Inventor UCKERT		•	First Given Name Stefan	Second Given Name
	Q.	Residence & Citizenship	City Garbsen	State or Foreign Country GERMANY	Country of Citizenship GERMANY
ľ	3	Mailing	Address	City	State & Zip Code/Country
		Address	Erich-Ollenhauer-Str. 3	Garbsen	D-30827 / GERMANY
	Signature of Inventor 203: Selec USel				07.01.2002
1	2	Full Name Of Inventor	Family Name JONAS	First Given Name Udo	Second Given Name
	0	Residence & Citizenship	City Hannover	State or Foreign Country GERMANY	Country of Citizenship GERMANY
	4	Mailing Address	Address Schopenhauer-Str. 27	City Hannover	State & Zip Code/Country D-30625 / GERMANY
	Signa	ture of Inventor 20	4:	Jan 7, 2002	
-					